

2010 (KUDA 012252). All verified by Charles W. Painter from photographs. New county records (Degenhardt et al. 1996. *Amphibians and Reptiles of New Mexico*. Univ. New Mexico Press, Albuquerque. xix + 431 pp.).

JAMES N. STUART, New Mexico Department of Game and Fish, Conservation Services Division, Santa Fe, New Mexico 87504-5112, USA (e-mail: james.stuart@state.nm.us); **MARK BJORKLUND**, 292 Aragon Avenue, Los Alamos, New Mexico 87544, USA (e-mail: mdbhb@q.com).

LITHOBATES HECKSCHERI (River Frog). USA: ALABAMA: BULLOCK Co.: The Wehle Tract (32.03151°N, 85.47390°W; WGS 84). 19 September 2011. Brian Folt. Verified by Craig Guyer. Auburn University Herpetological collection (photo voucher AHAP-D 326a,b). First county record and the first voucher for this species in Alabama since 1975 (Mount 1975. *The Reptiles and Amphibians of Alabama*. Auburn University Agricultural Experiment Station, Auburn, Alabama. vii + 347 pp.).

Found under log along shoreline of impoundment pond; captured by hand. Previously, *L. heckscheri* had been verified in only three localities in Alabama: one each in Mobile, Baldwin, and Escambia counties (Mount 1975, *op. cit.*). Whereas the historic Alabama localities are exclusively within the Lower Coastal Plains, this new record is farther north (171 km ENE from the nearest known location in Alabama), situated within the transitional zone between the Red Hills and Black Belt regions. Because this species is thought to be restricted to the Coastal Plains (Jensen et al. 2008. *Amphibians and Reptiles of Georgia*. University of Georgia Press, Athens. 575 pp.), this record is novel and suggests that other habitats might be suitable.

BRIAN P. FOLT, Auburn University, Department of Biological Sciences, 330 Funchess Hall, Auburn, Alabama 36849, USA; e-mail: brian.folt@gmail.com.

MELANOPHYRINISCUS KLAPPENBACHI (Klappenbach's Red-bellied Toad). BRAZIL, MATO GROSSO DO SUL, Municipality of Porto Murtinho, Fazenda Santo Antônio (21.527163°S, 57.832186°W, SAD 69); Fazenda Carandá (21.554553°S, 57.781036°W, SAD 69). 30 May 2011. I. B. Amaral, P. Landgraf-Filho, and D. I. Ferreira. Coleção Herpetológica do Museu de Ciências e Tecnologia PUCRS, Porto Alegre, RS, Brazil (MCP/PUC/RS 11944-11946, 11949-11969). Fazenda Patolá (21.686749°S, 57.719681°W, WGS 1984). 27 October 2008. F. L. Souza, M. Uetanabaro, and P. Landgraf-Filho. Coleção Zoológica de Referência da Universidade Federal de Mato Grosso do Sul, Campo Grande, MS, Brazil (ZUFMS AMP 1046). Verified by T. Grant. This species was previously known for Paraguay (southern Alto Paraguay, Presidente Hayes, and Ñeembucú Department: Brusquetti and Lavilla 2006. *Cuad. Herpetol.* 20[2]:1–79) and Argentina (Chaco, Formosa, and Northern Santa Fe and Santiago del Estero provinces: Baldo 2001. *Cuad. Herpetol.* 15[2]:141–142). First country record. The study site is located in the southern Pantanal, at the left bank of Paraguay River, the only Brazilian region encompassed by the Chaco biome. New localities extends know distribution 90 km N of Alto Paraguay, Paraguay, in Municipality of Porto Murtinho, Brazil.

IVAN BOREL AMARAL, Centro Nacional de Pesquisa e Conservação de Répteis e Anfíbios – RAN/ Instituto Chico Mendes de Conservação da Biodiversidade – ICMBio (e-mail: ivan.amaral@icmbio.gov.br); **PAULO LANDGRAF FILHO** (e-mail: paulograf@yahoo.com.br); **FRANCO LEANDRO SOUZA**, Centro de Ciências Biológicas e da Saúde, Universidade Federal de Mato Grosso do Sul, CEP 79070-900 Campo Grande, MS, Brazil (e-mail: franco.souza@ufms.br) and **MASAO UETANABARO** (e-mail: masao.uetanabaro@gmail.com).

PHLYCTIMANTIS LEONARDI (Olive Striped Frog). CAMEROON: EAST PROVINCE: Ngoum-Bandi (aka PK27), S border of Lobéké National Park (02.13881°N, 15.65567°E; WGS 84; 620 m elev.). 28 May 2010. Václav Gvoždík and Oldřich Kopecký. National Museum, Prague, Czech Republic. NMP6V 74437/1–5. Verified by Jean-Louis Amiet and Mark-Oliver Rödel. Previously recorded in Gabon, Equatorial Guinea (mainland), Republic of Congo, and western Democratic Republic of Congo (Schiotz et al. 2004. *In* IUCN 2011. *IUCN Red List of Threatened Species*, ver. 2011.1. <www.iucnredlist.org>). Four adult males (SVL 47.9–51.0 mm) and one adult female (SVL 54.8 mm) collected from small shallow pond at edge of primary forest. Males calling from grassy inundated banks and low bushes from dusk until ca. 2400 h. Pairs in amplexus and numerous tadpoles also observed. New record extends known range by ca. 50 km N from nearest localities in Republic of Congo. First species record for Cameroon and first record of genus from Cameroonian Congo Basin (cf. Amiet 2007. *Rev. Suisse Zool.* 114:87–126).

VÁCLAV GVOŽDÍK, Institute of Animal Physiology and Genetics, Academy of Sciences of the Czech Republic, Rumburska 89, CZ 277 21 Libečov & National Museum, Department of Zoology, Cirkusova 1740, CZ 193 00 Prague, Czech Republic (e-mail: vgvozdik@email.cz); **OLDŘICH KOPECKÝ**, Department of Zoology and Fisheries, Faculty of Agrobiological, Food and Natural Resources, Czech University of Life Sciences Prague, Kamycka 957, CZ 165 21 Prague, Czech Republic (e-mail: kopeckyo@af.czu.cz).

PSEUDACRIS CLARKII (Spotted Chorus Frog). USA: NEW MEXICO: QUAY Co.: Playa lake located on the south side of NM Hwy 231, 2.0 km W of intersection of NM Hwy 469 and NM Hwy 23; ca. 2.7 air km SW of Wheatland (34.89232°N, 103.37679°W, NAD1983; elev. 1440 m). 05 August 2011. Jessica A. Kissner. Verified by Toby Hibbitts. University of Kansas (KUDA digital images 012215–012218, and 012219 audio). First state record (Degenhardt et al. 1996. *Amphibians and Reptiles of New Mexico*. University of New Mexico Press, Albuquerque, New Mexico). Nearest previous record was at Muleshoe National Wildlife Refuge in Muleshoe, Texas, ca. 117 km airline SE from the new locality. At 2144 h, two adult males were heard and seen calling from the base of emergent vegetation after a 0.5 mm rainfall. Air temperature was 20.9°C and humidity was 83%, with cloudy skies and an average wind speed of 5.6 mph.

Field work was conducted under permit number 3318 issued by New Mexico Department of Game and Fish.

JESSICA A. KISSNER (e-mail: jessica.kissner@ttu.edu) and **KERRY L. GRIFFIS-KYLE**, Texas Tech University, Department of Natural Resources Management, Lubbock, Texas 79409, USA (e-mail: kerry.griffis-kyle@ttu.edu).

PSEUDACRIS MACULATA (Boreal Chorus Frog). USA: NEBRASKA: HARLAN Co.: Republican City (40.089455°N, 99.213149°W; elev. 612 m). 22 May 2011. Brian Hubbs. Verified by Dan Fogell. Natural History Museum of Los Angeles County photo voucher LACM PC 1563. New county record (Ballinger et al. 2010. *Amphibians and Reptiles of Nebraska*. Rusty Lizard Press, Oro Valley, Arizona. 400 pp.; Fogell 2010. *The Amphibians and Reptiles of Nebraska*. University of Nebraska Press, Lincoln, Nebraska. 158 pp.). Frog found in flooded roadside ditch.

BRIAN HUBBS, PO Box 26407, Tempe, Arizona 85285, USA; e-mail: tricolorbrian@hotmail.com.

RAORCHESTES PARVULUS (Karin Bubble-nest Frog). BANGLADESH: SYLHET DIVISION: Sylhet District: Khadimnagar National Park (24.940556°N, 91.93889°E; WGS 84; 46 m elev.). 29 April